

NCBI Protein

PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Books

Search **Protein** for

Limits Preview/Index History Clipboard Details

Display GenPept all to file

Range: from to Features: SNP CDD MGC HPRD STS

1: P77455. Reports MaoC protein (Phe...[gi:2494078] BLINK, Domains, Links

LOCUS P77455 681 aa linear BCT 13-SEP-2005
DEFINITION MaoC protein (Phenylacetic acid degradation protein paaZ).
ACCESSION P77455
VERSION P77455 GI:2494078
DBSOURCE swissprot: locus MAOC_ECOLI, accession [P77455](#);
 class: standard.
 extra accessions:O53009, created: Nov 1, 1997.
 sequence updated: Nov 1, 1997.
 annotation updated: Sep 13, 2005.
 xrefs: [X97452.1](#), [CAA66089.1](#), [U00096.2](#), [AAC74469.1](#), [D90777.1](#),
[BAA14997.1](#), [F64889](#)
 xrefs (non-sequence databases): EchoBASEB3498, EcoGeneEG13735,
 InterProIPR002086, InterProIPR002539, InterProIPR011966,
 PfamPF00171, PfamPF01575, TIGRFAMsTIGR02278, PROSITEPS00070,
 PROSITEPS00687
KEYWORDS Complete proteome; Oxidoreductase.
SOURCE Escherichia coli
ORGANISM [Escherichia coli](#)
 Bacteria; Proteobacteria; Gammaproteobacteria; Enterobacteriales;
 Enterobacteriaceae; Escherichia.
REFERENCE 1 (residues 1 to 681)
AUTHORS Fernandez,A., Minambres,B., Garcia,B., Olivera,E.R., Luengo,J.M.,
 Garcia,J.L. and Diaz,E.
TITLE Catabolism of phenylacetic acid in Escherichia coli.
 Characterization of a new aerobic hybrid pathway
JOURNAL J. Biol. Chem. 273 (40), 25974-25986 (1998)
PUBMED [9748275](#)
REMARK NUCLEOTIDE SEQUENCE.
 STRAIN=W / ATCC 11105
REFERENCE 2 (residues 1 to 681)
AUTHORS Blattner,F.R., Plunkett,G. III, Bloch,C.A., Perna,N.T., Burland,V.,
 Riley,M., Collado-Vides,J., Glasner,J.D., Rode,C.K., Mayhew,G.F.,
 Gregor,J., Davis,N.W., Kirkpatrick,H.A., Goeden,M.A., Rose,D.J.,
 Mau,B. and Shao,Y.
TITLE The complete genome sequence of Escherichia coli K-12
JOURNAL Science 277 (5331), 1453-1474 (1997)
PUBMED [9278503](#)
REMARK NUCLEOTIDE SEQUENCE [LARGE SCALE GENOMIC DNA].
 STRAIN=K12 / MG1655
REFERENCE 3 (residues 1 to 681)
AUTHORS Aiba,H., Baba,T., Fujita,K., Hayashi,K., Inada,T., Isono,K.,
 Itoh,T., Kasai,H., Kashimoto,K., Kimura,S., Kitakawa,M.,
 Kitagawa,M., Makino,K., Miki,T., Mizobuchi,K., Mori,H., Mori,T.,
 Motomura,K., Nakade,S., Nakamura,Y., Nashimoto,H., Nishio,Y.,
 Oshima,T., Saito,N., Sampei,G., Seki,Y., Sivasundaram,S.,

Tagami,H., Takeda,J., Takemoto,K., Takeuchi,Y., Wada,C.,
 Yamamoto,Y. and Horiuchi,T.
TITLE A 570-kb DNA sequence of the Escherichia coli K-12 genome
 corresponding to the 28.0-40.1 min region on the linkage map
JOURNAL DNA Res. 3 (6), 363-377 (1996)
PUBMED [9097039](#)
REMARK NUCLEOTIDE SEQUENCE [LARGE SCALE GENOMIC DNA].
 STRAIN=K12
REFERENCE 4 (residues 1 to 681)
AUTHORS Ferrandez,A., Garcia,J.L. and Diaz,E.
TITLE Transcriptional regulation of the divergent paa catabolic operons
 for phenylacetic acid degradation in Escherichia coli
JOURNAL J. Biol. Chem. 275 (16), 12214-12222 (2000)
PUBMED [10766858](#)
REMARK TRANSCRIPTIONAL REGULATION.
COMMENT On May 27, 2005 this sequence version replaced gi:[7466625](#).
 [PATHWAY] Phenylacetic acid aerobic catabolism.
 [INDUCTION] Activated by cAMP receptor protein (CRP) and
 integration host factor (IHF). Inhibited by paaX.
 [SIMILARITY] Belongs to the aldehyde dehydrogenase family.
FEATURES Location/Qualifiers
source 1..681
 /organism="Escherichia coli"
 /db_xref="taxon:[562](#)"
gene 1..681
 /gene="maoC"
 /locus_tag="b1387"
 /note="synonym: paaZ"
Protein 1..681
 /gene="maoC"
 /locus_tag="b1387"
 /product="MaoC protein"
Region 143
 /gene="maoC"
 /locus_tag="b1387"
 /region_name="Variant"
 /note="L -> V (in strain W)."
 /evidence=experimental
Site 256
 /gene="maoC"
 /locus_tag="b1387"
 /site_type="active"
 /note="By similarity."
 /evidence=not_experimental
Site 295
 /gene="maoC"
 /locus_tag="b1387"
 /site_type="active"
 /note="By similarity."
 /evidence=not_experimental
Region 440
 /gene="maoC"
 /locus_tag="b1387"
 /region_name="Variant"
 /note="A -> T (in strain W)."
 /evidence=experimental
Region 611
 /gene="maoC"
 /locus_tag="b1387"
 /region_name="Variant"

/note="S -> N (in strain W)." /evidence=experimental

ORIGIN

1 mqqlasflsg twqsgrgrsr lihhaisgea lwevtsegld maaarqfaie kgapalramt
61 fieraamlka vakhllseke rfyalsaqtg atradswdi eggigtlfty aslgsrelpd
121 dtlwpedeli plskeggfaa rhlltsksgv avhinafnfp cwgmleklap twlggmpaii
181 kpatataqlt qamvksivds glvpegaisl icgsagdlld hldsqdvvtf tgsaatggml
241 rvqpnnivaks ipftmeadsl nccvlgedvt pdqpefalfi revvremttk agqkctairr
301 iivpqalvna vsdalvarlq kvvvgdpae gvkmgalvna eqradvqekv nillaagcei
361 rlbggqadlsa agaffpptll ycpqpdetpa vhateafgpv atlmpaqnqr halqlacagg
421 gslagtlvta dpqiarqfia daarthgriq ilneesakes tghgsplpql vhggpgragg
481 geelggrrav khymqravq gsptmlaais kqwvrgakve edrihpfrky feelqpgds1
541 ltprrtmtea divnfaclsg dhfyahmdki aaaesifger vvhgyfvlsa aaglfvdagv
601 gpvianygle slrfiepvkp gdtiqvrllc krktlkqrs aekptgvve wavevfnqhq
661 tpvalysilt lvarqhgdfv d

//

[Disclaimer](#) | [Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)

Feb 9 2005 14:31:10

[ExPASy Home page](#)[Site Map](#)[Search ExPASy](#)[Contact us](#)[Swiss-Prot](#)Search for

UniProtKB/Swiss-Prot entry P77455

[Printer-friendly view](#)[Submit update](#)[Quick Blast!](#)
[\[Entry info\]](#)
[\[Name and origin\]](#)
[\[References\]](#)
[\[Comments\]](#)
[\[Cross-references\]](#)
[\[Keywords\]](#)

[\[Features\]](#)
[\[Sequence\]](#)
[\[Tools\]](#)

Note: most headings are clickable, even if they don't appear as links. They link to the user manual or other documents.

Entry information

Entry name	MAOC_ECOLI
Primary accession number	P77455
Secondary accession number	O53009
Entered in Swiss-Prot in	Release 35, November 1997
Sequence was last modified in	Release 35, November 1997
Annotations were last modified in	Release 48, September 2005

Name and origin of the protein

Protein name	MaoC protein
Synonym	Phenylacetic acid degradation protein paaZ
Gene name	Name: maoC

From

Taxonomy
Escherichia coli [TaxID: 562]
Bacteria; Proteobacteria; Gammaproteobacteria; Enterobacteriales;
Enterobacteriaceae; Escherichia.

References

[1] NUCLEOTIDE SEQUENCE.

STRAIN=W / ATCC 11105;
DOI=10.1074/jbc.273.40.25974;MEDLINE=98421522;PubMed=9748275 [NCBI, ExPASy, EBI, Israel, Japan]
Ferrandez A., Minambres B., Garcia B., Olivera E.R., Luengo J.M., Garcia J.L., Diaz E.;
"Catabolism of phenylacetic acid in Escherichia coli. Characterization of a new aerobic hybrid pathway.";
J. Biol. Chem. 273:25974-25986(1998).

[2] NUCLEOTIDE SEQUENCE [LARGE SCALE GENOMIC DNA].

STRAIN=K12 / MG1655;
DOI=10.1126/science.277.5331.1453;MEDLINE=97426617;PubMed=9278503 [NCBI, ExPASy, EBI, Israel, Japan]
Blattner F.R., Plunkett G. III, Bloch C.A., Perna N.T., Burland V., Riley M., Collado-Vides J., Glasner J.D., Rode C.K., Mayhew G.F., Gregor J., Davis N.W., Kirkpatrick H.A., Goeden M.A., Rose D.J., Mau B., Shao Y.;
"The complete genome sequence of Escherichia coli K-12.";
Science 277:1453-1474(1997).

[3]

NUCLEOTIDE SEQUENCE [LARGE SCALE GENOMIC DNA].

STRAIN=K12;

MEDLINE=97251357;PubMed=9097039 [NCBI, ExPASy, EBI, Israel, Japan]

Aiba H., Baba T., Fujita K., Hayashi K., Inada T., Isono K., Itoh T., Kasai H., Kashimoto K., Kimura S., Kitakawa M., Kitagawa M., Makino K., Miki T., Mizobuchi K., Mori H., Mori T., Motomura K., Nakade S., Horiuchi T.;

"A 570-kb DNA sequence of the Escherichia coli K-12 genome corresponding to the 28.0-40.1 min region on the linkage map.";

DNA Res. 3:363-377(1996).

[4] TRANSCRIPTIONAL REGULATION.

DOI=10.1074/jbc.275.16.12214;MEDLINE=20229831;PubMed=10766858 [NCBI, ExPASy, EBI, Israel, Japan]

Ferrandez A., Garcia J.L., Diaz E.;

"Transcriptional regulation of the divergent paa catabolic operons for phenylacetic acid degradation in Escherichia coli.";

J. Biol. Chem. 275:12214-12222(2000).

Comments

- **PATHWAY:** Phenylacetic acid aerobic catabolism.
- **INDUCTION:** Activated by cAMP receptor protein (CRP) and integration host factor (IHF). Inhibited by paaX.
- **SIMILARITY:** Belongs to the aldehyde dehydrogenase family.

Copyright

This Swiss-Prot entry is copyright. It is produced through a collaboration between the Swiss Institute of Bioinformatics and the EMBL outstation - the European Bioinformatics Institute. There are no restrictions on its use as long as its content is in no way modified and this statement is not removed.


Cross-references

	X97452; CAA66089.1; -;	[EMBL / GenBank / DDBJ]
	Genomic_DNA.	[CoDingSequence]
EMBL	U00096; AAC74469.1; -;	[EMBL / GenBank / DDBJ]
	Genomic_DNA.	[CoDingSequence]
	D90777; BAA14997.1; -;	[EMBL / GenBank / DDBJ]
	Genomic_DNA.	[CoDingSequence]
PIR	F64889; F64889.	
EchoBASE	EB3498; -.	
EcoGene	EG13735; maoC.	
EcoCyc	EG13735; maoC.	
CMR	P77455; b1387.	
	IPR002086; Aldehyd_dehydrog.	
InterPro	IPR002539; MaoC_dehydratas.	
	IPR011966; PaaN-DH.	
	Graphical view of domain structure.	
Pfam	PF00171; Aldedh; 1.	
	PF01575; MaoC_dehydratas; 1.	
	Pfam graphical view of domain structure.	
TIGRFAMs	TIGR02278; PaaN-DH; 1.	
PROSITE	PS00070; ALDEHYDE_DEHYDR_CYS; FALSE_NEG.	
	PS00687; ALDEHYDE_DEHYDR_GLU; FALSE_NEG.	
ProDom	[Domain structure / List of seq. sharing at least 1 domain]	
HOGENOM	[Family / Alignment / Tree]	
BLOCKS	P77455.	

ProtoNet P77455.
 ProtoMap P77455.
 PRESAGE P77455.
 DIP P77455.
 ModBase P77455.
 SWISS-2DPAGE Get region on 2D PAGE.
 UniRef View cluster of proteins with at least 50% / 90% identity.
Keywords

Complete proteome; Oxidoreductase.

Features



Feature table viewer

Key	From	To	Length	Description
ACT_SITE	256	256	1	By similarity.
ACT_SITE	295	295	1	By similarity.
VARIANT	143	143	1	L → V (in strain W).
VARIANT	440	440	1	A → T (in strain W).
VARIANT	611	611	1	S → N (in strain W).

Sequence information

Length: 681 Molecular weight: 73003 CRC64: A631B97AA37A1C3E [This is a checksum on the AA sequence]

10	20	30	40	50	60
MQQLASFLSG	TWQSGRGRSR	LIHHAIISGEA	LWEVTSEG LD	MAAARQFAIE	KGAPALRAMT
70	80	90	100	110	120
FIERAAMLKA	VAKHLLSEKE	RFYALSAQTG	ATRADSWVDI	EGGIGTLFTY	ASLGSRELPD
130	140	150	160	170	180
DTLWPEDELI	PLSKEGGFAA	RHLLTSKSGV	AVHINAFNFP	CWGMLEKLAP	TWLGGMPAI
190	200	210	220	230	240
KPATATAQLT	QAMVKSIVDS	GLVPEGAISL	ICGSAGDL	LD HLDSQDVVT	TF TGSAATGQML
250	260	270	280	290	300
RVQPNI VAKS	IPFTMEADSL	NCCVLGEDVT	PDQPEFALFI	REVVREM	TTK AGQKCTAIRR
310	320	330	340	350	360
IIVPQALVNA	VSDALVARLQ	KVVVGDP	AE GVKGALVNA	EQRADVQE KV	NILLAAGCE I
370	380	390	400	410	420
RLGGQADLSA	AGAFFPPTILL	YCPQPDET PA	VHATEAFGPV	ATLMPAQNQR	HALQLACAGG
430	440	450	460	470	480
GSLAGTLVTA	DPQIARQFIA	DAARTHGRIQ	ILNEESAKES	TGHGSPLPQL	VHGPGRAGG
490	500	510	520	530	540
GEELGGGLRAV	KHYM QRTAVQ	GSPTMLAAIS	KQWVRGAKVE	EDRIHPFRKY	FEELQPGDSL
550	560	570	580	590	600
LTPRRTMTEA	DIVNFACLSG	DHFYAHMDKI	AAAESIFGER	VVHGYFVLSA	AAGLFVDAGV

610 620 630 640 650 660
GPVIANYGLE SLRFIEPVKP GDTIQVRLTC KRKTLKKQRS AEEKPTGVVE WAVEVFNQHQ
670 680
TPVALYSILT LVARQHGDFV D

P77455 in FASTA
format

[View entry in original UniProtKB/Swiss-Prot format](#)

[View entry in raw text format \(no links\)](#)

[Report form for errors/updates in this UniProtKB/Swiss-Prot entry](#)

BLAST BLAST submission on
ExPASy/SIB
or at NCBI (USA)



Sequence analysis tools: ProtParam, ProtScale,
Compute pI/Mw, PeptideMass, PeptideCutter,
Dotlet (Java)



ScanProsite, MotifScan



Submit a homology modeling request to SWISS-
MODEL

[ExPASy Home page](#)

[Site Map](#)

[Search ExPASy](#)

[Contact us](#)

[Swiss-Prot](#)

Hosted by NCSC
US

Mirror
sites:

Australia Bolivia Brazil Canada Korea Switzerland Taiwan